

NUTRITIONAL VALUE AND PHYSICOCHEMICAL COMPOSITION OF PEARL MILLET (*Pennisetum glaucum*) PRODUCED IN BENIN

Kifouli Adéoti^{*1}, Sonagnon H.S. Kouhoundé¹, Pacôme A. Noumavo^{1,2}, Farid Baba-Moussa¹, Fatiou Toukourou¹

Address(es):

¹Laboratoire de Microbiologie et des Technologies Alimentaires, Champ de Foire ISBA, Faculté des Sciences et Techniques, Université d'Abomey-Calavi, Bénin.

²Laboratoire de Biologie et de Typage Moléculaire en Microbiologie, Campus d'Abomey-Calavi, Faculté des Sciences et Techniques, Université d'Abomey-Calavi, Bénin.

*Corresponding author: zoulade@yahoo.fr

doi: 10.15414/jmbfs.2017.7.1.92-96

ARTICLE INFO

Received 15. 11. 2016
Revised 30. 6. 2017
Accepted 20. 7. 2017
Published 1. 8. 2017

Regular article



ABSTRACT

The high prices of animal foods and limited income earned in developing countries have resulted in their dependency on cereal-based preparations as staple food. Cereals such as maize, sorghum or pearl millet are often used in the production of various traditional foods and beverages in many African countries including Benin. In the republic of Benin, the nutritive value of cereals such as maize and sorghum is well documented. However, the nutritional value of pearl millet varieties produced in Benin remains to be investigated. The aim of this study was to assess the nutritional value of 22 varieties of pearl millet produced in Benin. After samples collection, the pearl millet grains were milled into fine powder and their compositions in six minerals (Na^+ , K^+ , Ca^{2+} , Mg^{2+} , Fe^{2+} , Zn^{2+}) and proximate (ash, dry matter, starch, protein, total and reducing sugars) were determined using standard analytical AOAC methods. Next, we assessed the relationship between these variables using the Pearson Correlation Analysis (PCA). We found that protein, total and reducing sugars levels varied widely (ranged from 1.86 to 93.4 mg/g, 178.64 to 652.54 mg/g and 16.62 to 174.22 mg/g, respectively). Additionally, we found a highly significant correlation ($P < 0.001$) between levels of starch and amylose. Mineral levels also varied widely, with some millet cultivars being particularly enriched in iron and magnesium (levels ranged from 13.85 to 2766.31 mg/kg, and from 340.27 to 4769.9 mg/kg, respectively). Four groups of pearl millet can be distinguished based on data from the PCA: iron-rich millets (group G1), carbohydrate-rich millets (group G2), and two less nutritious millets (groups G3 and G4). This study opens new avenues for millet fortification and provides opportunity to increase farmers' awareness in selecting pearl millet varieties for reducing malnutrition.

Keywords: Pearl millet, nutritional value, physicochemical composition, food security, Benin

INTRODUCTION

Africa is the center of origin and a major producer of several cereals (Macauley, 2015). Cereals are the major dietary energy supplies and are consumed in a variety of form depending on the ethnic or religious affiliation (Idem and Showemimo, 2004). The important cultivated cereals crops in Africa are maize, millet, rice, sorghum and wheat (Macauley, 2015). Pearl millet (*Pennisetum glaucum*), a hardy cereal crop compared with wheat (*Triticum aestivum*) and rice (*Oryza sativa* L.), is grown in regions characterized by relatively low rainfall owing to its ability to tolerate and survive under continuous or intermittent drought conditions (Jain and Bal, 1997). It is a drought tolerant cereal crop grown primarily as a food grain in India and Africa (Dykes and Rooney, 2007; Hoover et al., 1996). Because of its good performances in low-fertility soils such as semi-arid regions of Africa and Southeast Asia, Pearl millet is an important staple cereal cultivated in regions severely affected by malnutrition. It has a high nutritional value compared to others cereals and presents therefore a high potential in contributing to food and nutrition security (Shweta, 2015). The cultivated area of millet on the African continent was estimated to 19,998,008 hectares, with annual production reaching 16,008,838 tons (FAO, 2015; Macauley and Ramadjita, 2015). Pearl millet production ranked third among cereals in Africa. In Benin, the national production (27000 tons in 2013, USDA 2013) ranks fourth among cereals after rice, maize and sorghum. Importantly, its production ranks second in the Northern part of the country, and could therefore be a potential target for studying diversity of millets produced within Benin. Being a drought-tolerant crop, pearl millet is cultivated in the dry Northern regions of Benin under increasingly short and marginal raining seasons. It is used in various local food preparations like local porridge, cakes and traditional beverages.

Previous studies showed that millet has several potential health benefits, partly attributed to its polyphenol and dietary fiber contents (Devi et al., 2014). Depending on the country, millet is sometimes used directly in preparations

(Adebiyi et al., 2016) or mixed with other grains like sorghum or cowpea in order to improve the nutritional quality of food (Modu et al., 2010; Almeida-Dominguez et al., 1993;). In Burkina-Faso for example, it is used as complementary food for young children when pearl millet is fermented (Songré-Ouattara et al., 2009, 2008; Tou et al., 2006). The importance of this crop for developing countries incited biotechnologists to converge their attention on the improvement of some species of millet (Kothari et al., 2005). On the other hand, epidemiological studies have demonstrated that regular consumption of whole grain cereals and their products can protect against the risk of cardiovascular diseases, type II diabetes, gastrointestinal cancers and a range of other disorders (McKeown et al., 2002).

Although many studies have been conducted on the plants and grains, the assessment of the nutritional factors of pearl millet varieties produced in Benin has not been performed. Furthermore, pearl millet varieties cultivated in Benin are not well documented. We devised this study to understand to what extent intake of pearl millets could contribute to food security in the Northern Benin. The aim of our study was to determine concentrations of six selected minerals (iron, zinc, magnesium, calcium, potassium) and proximate compositions of 22 varieties of pearl millet produced in Benin.

MATERIAL AND METHODS

Samples collection

A total of 22 cultivars of pearl millets were collected from two regions of Northern part of Benin Republic (Atacora-Donga and Borgou-Alibori). Following collection, the grains were washed thoroughly to remove any debris or dust particles, milled into fine powder using house blender. Table 1 presents the cultivars description along with their vernacular names and the names of the sites (villages) of provenance.

Table 1 List of pearl millet samples collected

Samples order	Region of Benin	Vernacular name	Collecting site	Samples code
1		<i>Eyémata</i>	Manta (kouwinkou)	CA
2		<i>Amala korolomè</i>	Sosso II	CB
3		<i>Sowariya</i>	Boribansifa	CC
4		<i>Zokpéra</i>	Tanéka-koko	CD
5		<i>Eyomata</i>	Pam-Pam	CE
6		<i>Amala koukpètè</i>	Katanga	CF
7		<i>Zokpéra</i>	Barèi	CG
8	Atacora-Donga	<i>Diyématrìe</i>	Kouwotchirgou	CH
9		<i>Sowariya</i>	Yinrèboudé	CI
10		<i>Amala Kouhouloumè</i>	Sosso II	CJ
11		<i>Amala koukpètè</i>	Sosso II	CK
12		<i>Yomaga</i>	Tampatou	CL
13		<i>Iyaku</i>	Goun-Goun	CM
14		<i>Hinnitchiré</i>	Karimama centre	CN
15		<i>Guero</i>	Diapéou	CO
16		<i>Idié</i>	Goumori	CP
17		<i>Wea</i>	Kalalé centre	CQ
18		<i>Eyomata</i>	Boérou	CR
19	Borgou-Alibori	<i>Meïwaré</i>	Wanrarou	CS
20		<i>Gbéi</i>	Sinendé centre	CT
21		<i>Sonna</i>	Bodjekali	CU
22		<i>Iyasé</i>	Goun-Goun	CV

Proximate composition analysis

Proximate composition analysis consists of evaluation of proteins, total and reducing sugar, starch and amylose content according to the standard procedures of official method of analysis of the Association of Official Analytical Chemists (AOAC, 2000). The protein (total nitrogen) content was determined by Kjeldahl method using 6.25 as conversion constant. The total sugars were measured by spectrometry method at 492 nm using D-glucose as standard. Reducing sugars were also measured by spectrometry (Thermo Scientific Spectrophotometer, Biomate 3S, number 335904 series, 2004) method with a reagent 3.5 dinitrosalicylic acid (DNS) at 546 nm using maltose as standard. Starch content was determined by reading solution floated after preparing the (hydrolysis of starch) medium using the different pearl millet flour sample, sodium hydroxide (1 M) and chlorhydric acid (1 M) at 580 nm using BioMate™ 3S Spectrophotometer.

Mineral composition analysis

The minerals such as calcium were determined from aliquots of the solutions obtained from ash by established flame Atomic Absorption Spectrophotometry procedures (mineralization, calibration and dosage, 2003) as described by

Kouassi et al. (2013). Solution was prepared for each sample of variety. The sodium (Na⁺) and potassium (K⁺) were determined using flame photometer following the method previously described by Oboh and Oladunmoye (2007).

Statistical analysis

The results obtained were analyzed using descriptive statistics (means and standard deviation). Analysis of variance, Pearson correlation analysis and Principal Component Analysis (correlation matrix were generated to assess the correlation) were carried out using Statistical Package of Social Sciences (SPSS version 20.0).

RESULTS AND DISCUSSION

Proximate composition

The proximate composition analysis including the protein, total sugars, reducing sugars, starch and amylose content was assessed for the 22 cultivars of pearl millet. Ca²⁺, magnesium (Mg²⁺), iron (Fe²⁺) and zinc (Zn²⁺) The obtained results are presented in Table 2 below.

Table 2 Proteins, sugars, starch, and amylose compositions in selected cultivars of pearl millets

Samples code	Parameters (Mean ± SD)				
	Proteins	Total sugar	Reducing sugar	Starch	Amylose
CA	3.41±1.55	178.64±7.88	16.62±4.30	50.51±7.89	49.67±2.86
CB	26.68±1.55	219.50±3.17	40.48±2.69	201.00±5.17	124.15±3.38
CC	12.33±2.32	303.45±4.50	33.81±4.30	263.86±1.38	211.52±5.17
CD	14.66±2.33	394.87±5.74	61.55±2.92	302.89±7.70	247.38±8.17
CE	26.68±1.55	429.91±2.30	144.15±1.46	196.11±8.30	130.85±3.82
CF	18.54±0.78	514.78±1.50	41.69±5.73	510.63±12.78	375.71±19.84
CG	45.69±3.89	579.91±2.63	63.90±1.43	452.66±5.67	412.90±3.58
CH	15.82±3.10	421.00±2.57	173.47±7.02	101.16±8.47	65.51±0.156
CI	8.07±1.55	377.05±10.72	47.66±1.46	372.84±6.05	282.01±10.14
CJ	17.00±2.33	594.62±5.36	77.63±2.92	507.59±9.81	492.73±13.66
CK	1.86±1.55	195.58±2.57	59.79±1.403	416.58±11.44	392.60±17.91
CL	10.78±2.32	337.98±3.75	174.22±12.89	163.67±9.05	99.42±6.72
CM	43.75±1.55	413.24±1.88	57.45±2.86	445.15±22.77	429.17±10.19
CN	9.62±3.10	495.64±6.76	33.81±4.30	299.40±0.97	274.31±5.57
CO	66.63±0.78	605.77±2.94	40.14±4.21	76.07±2.50	39.78±2.08
CP	15.82±1.55	652.54±2.25	46.00±2.86	146.13±9.93	139.03±3.38
CQ	20.48±1.55	580.85±2.25	57.45±2.86	383.68±6.50	350.00±9.02
CR	7.29±1.55	488.28±4.78	78.74±2.81	570.07±24.76	479.56±14.21
CS	65.08±0.77	465.85±3.309	119.44±2.81	515.295±11.07	491.40±15.79
CT	93.40±1.55	359.05±3.06	85.67±4.38	478.35±6.00	342.78±12.10
CU	8.84±1.55	392.41±3.00	129.09±2.86	443.90±1.65	347.67±5.094
CV	32.11±3.10	604.77±4.21	79.82±4.38	560.55±7.81	390.44±25.734

The analysis of this table reveals that the protein contents vary from 1.86 mg/g (CK) to 93.4 mg/g (CT). This result shows that some of varieties have very low protein content. Among the 22 studied cultivars, six have a protein content less than 10 mg/g. However, the protein content obtained in this study is within the range of values reported in a study conducted in South Eastern USA (Yadav et al., 2014). These samples were found in both departments. Several studies showed that millet is a cereal relatively poor in protein (Dykes and Rooney, 2007; Vanisha et al., 2011; Shweta, 2015). The total sugar content varied from 178.64 mg/g (CA) to 652.54 (CP) mg/g while the reducing sugar content varied from 16.62 mg/g (CA) to 174.22 mg/g (CL). Among the nutritional benefits, the hypoglycemic property made millets particularly popular for their use in the management of type II diabetes. This hypoglycemic characteristic could be attributed to the presence of polyphenols, soluble fiber, and low level of reducing sugars. Consistent with this, we found low levels of sugars in the millet cultivars under study. Similar finding was reported by Taylor et al. (2006). For the starch content, its values vary from 50.51 mg/g (CA) to 570.07 mg/g (CR). However, the highest value obtained for amylose content was 491.40 mg/g (CS). Starch contents were significantly correlated with amylose levels (p <0.001). The presence of a large amount of starch in millet could decrease its digestibility. Indeed, Rooney and Pflugfeder (1986) showed that digestibility of starch is inversely proportional to amylose and amylopectin contents. Similar observation has been reported in previous studies (Laurent, 1988; Michalet-Dorea et Sauvante, 1989) of others cereals such as maize, sorghum, wheat, barley. However, the lower the proportion of amylose, the more gelatinization and vice

versa. More the starch resist to gelatinization, more it was converted by alpha-amylases to glucose, which tends to raise the glycemic index (Lopez-Rubio et al., 2009; Zhu et al., 2011).

Nevertheless by the fermentation, hydrolysis of millet carbohydrates not only increases digestibility but also its nutritional quality. This justifies its use in some countries to supplement the diets of young children (Songré-Ouattara et al., 2008). This quality improvement is the result of a cascade enzymatic reactions performed by the fermentation flora and mainly lactic acid bacteria (Songré-Ouattara et al., 2009).

Significant differences were observed in the proximate composition of millet samples under study. These differences were very highly significant for protein and sugar contents. It was noted that the samples collected from Borgou-Alibori region presented a best nutritional value than the samples collected from the region of Atacora-Donga. Similar findings have been reported in sorghum and millet (Almeida-Dominguez et al., 1991). These changes in the nutritional composition of the grains were found to be associated with both genetic and environmental factors (temperature, pluviometry, light intensity, physico-chemical characteristics of soil, etc.)

Mineral composition

The main minerals sought in pearl millet samples were Ca²⁺, Mg²⁺, K⁺, Na⁺, Fe²⁺ and Zn²⁺. In addition to these inorganic elements, the dry matter and ash were measured and the results are presented in Table 3.

Table 3 Micronutrient composition of flour obtained from selected cultivar of pearl millet

Samples code	Parameters (mg/kg)							
	% DM	Ash	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	Fe ²⁺	Zn ²⁺
CA	87.87	1.66	547.64	340.27	4401	196.19	36.07	481.03
CB	88.26	1.73	344.42	2775.18	3492	84.02	127.01	164.38
CC	88.01	1.54	424.36	2263.10	3824	240.47	70.92	214.14
CD	87.52	1.21	391.01	2081.72	3087	101.65	32.84	809.38
CE	88.72	2.9	94.02	1697.08	4128	78.04	1670.71	483.83
CF	88.31	1.36	187.50	2675.23	3648	85.58	37.69	269.53
CG	93.4	1.45	231.66	1588.41	3457	92.49	115.63	148.64
CH	87.98	1.60	75.61	1829.86	3920	105.06	63.13	313.79
CI	88.31	2.72	47.47	1646.84	3041	186.06	919.68	363.99
CJ	88.84	1.55	31.77	1566.87	3341	78.28	148.36	333.66
CK	87.37	2.55	88.56	1330.55	3425	106.28	248.89	547.97
CL	88.12	2.39	204.92	2689.41	3409	88.47	2766.31	528.07
CM	87.88	1.62	47.29	1544.98	3868	89.38	78.42	43.34
CN	88.57	1.36	219.60	2382.03	3319	68.31	37.25	56.86
CO	89.05	1.58	77.97	2264.32	3226	67.74	132.16	187.13
CP	88.21	1.45	676.69	2274.43	3022	58.37	39.47	233.08
CQ	88.84	2.12	500	2126.66	3352	114.21	103.97	200.75
CR	88.38	1.09	377.95	2772.51	3557	233.99	156.69	155.51
CS	89.35	1.56	576	1829.52	3663	132.28	56	204
CT	88.72	1.76	728.71	2482.29	4050	134.11	51.48	73.26
CU	87.71	1.53	468.29	4769.17	4082	179.95	13.85	261.46
CV	88.39	1.20	339.44	4415.27	3457	121.25	318.61	246.86

DM = Dry Matter; Ca²⁺ = Calcium; Mg²⁺ = Magnesium; K⁺ = Potassium; Na⁺ = Sodium; Fe²⁺ = Iron; Zn²⁺ = Zinc.

As shown in Table 3 above, the studied cultivars of pearl millet are rich in minerals and contain more minerals than corn and sorghum (Almeida-Dominguez et al., 1993, 1991; N’guessan et al., 2014). An analysis of this table reveals that the Ca²⁺ content in the analyzed samples varied between 31.77 mg/kg (CJ) and 728.71 mg/kg (CT). The levels of Mg²⁺ and Na⁺ ranged from 340.27 mg/kg (CA) to 4769.9 mg/kg (CU), and 58.37 mg/kg (CP) to 240.47 mg/kg (CC), respectively. For Fe²⁺ and Zn²⁺ the levels ranged from 13.85 mg/kg (CU) to 2766.31 mg/kg (CL), and 43.34 mg/kg (CM) to 809.38 mg/kg (CD), respectively. Indeed millet cultivated in Benin is an important source of Fe²⁺, K⁺, Mg²⁺ and Ca²⁺. Note that for Ca²⁺ and Mg²⁺ the highest levels were obtained in samples collected from the region of Atacora-Donga. On the other hand, the highest levels of K⁺, Na⁺, Fe²⁺ and Zn²⁺ were recorded in samples collected from the Borgou-Alibori region. No significant differences were observed for dry matter, ash and K⁺, unlike all other minerals analyzed. Nevertheless, Mouquet-Rivier et al. (2008) reported lower content of iron and zinc for millet produced in Burkina Faso. Apart from all other minerals, it was noted that this cereal contains a very

low content of Ca²⁺. According to Recommended Dietary Allowance (RDA), this cereal cannot cover alone needs of Ca²⁺ for young children (which between 1,000 and 2,500 mg for children aged ≤ 8 years). Flour millet is also used in the feeding of young children in Benin. In Nigeria, Akeredolu et al. (2005) reported that this cereal can be used in dietary supplement to promote growth of children because of its nutrient content.

Correlation among nutritional parameters

The pearl millet and sorghum have the advantage of serving in both the nutrition of young children and adult ration where the antioxidants are very sought (Obizoba and Atii, 1994; Ragae et al., 2006). This characteristic depends on the mineral and nutritional composition. In this section, the correlations between variables were analyzed according to Pearson method. The table showed below presents the correlations between determined parameters (Table 4).

Table 4 Pearson correlation matrices between the physicochemical and nutritional parameters of pearl millet collected in Benin

Parameters	DM	Ash	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	Fe ²⁺	Zn ²⁺	Protein	Total sugars	Reducing sugars	Starch
Ash	-0.114											
Ca ²⁺	-0.030	-0.333										
Mg ²⁺	-0.144	-0.324	0.211									
K ⁺	-0.112	0.049	0.187	-0.063								
Na ⁺	-0.190	-0.083	0.313	0.080	0.362*							
Fe ²⁺	-0.054	0.655***	-0.302	0.007	-0.036	-0.134						
Zn ²⁺	-0.345	0.343	-0.154	-0.230	-0.098	-0.018	0.395*					
Protein	0.411*	-0.123	0.236	0.024	0.128	-0.234	-0.163	-0.460*				
Total sugar	0.438*	-0.369*	-0.036	0.256	-0.453*	-0.388*	-0.157	-0.353	0.243			
Reducing sugar	-0.029	-0.131	-0.131	0.247	0.268	-0.097	0.554**	0.213	0.056	0.005		
Starch	0.201	0.047	0.047	0.333	-0.073	0.192	-0.253	-0.288	0.181	0.267	-0.022	
Amylose	0.242	0.019	0.019	0.158	-0.104	0.149	-0.312	-0.287	0.144	0.278	-0.065	0.963***

DM: dry matter; Ca²⁺: calcium; Mg²⁺: magnesium; K⁺: potassium; Na⁺: sodium; Zn²⁺: zinc; Fe²⁺: Iron; * indicate a significant correlation (p < 0.05); ** indicate a highly significant correlation (p < 0.01); *** indicate a very highly significant correlation (p < 0.001)

It is clear from this analysis that the low ash content induces the high protein content. For sugar content a significant correlation was also observed with the ash (p < 0.05). This observation is in agreement with an earlier study on grain of sorghum (Subramanian and Jambunathan, 1982). However, some authors have observed an inverse association between the protein content and starch content of grains. This relationship had already been demonstrated in sorghum by a group of researchers (Chandrashekar and Kirleis, 1988). The previous authors showed that the presence of proteins affects the starch gelatinization. The gelatinization was also influenced by the amylose content. Our study showed highly significant correlation (p < 0.001) between amylose and starch. The study conducted on rice showed that, higher is the proportion of amylose, the lower is the gelatinization and vice versa (Chung et al., 2011). In another study, it was demonstrated that the differences between the physicochemical and nutritional parameters of starch are also a function of the structural characteristics of cultivars (Chávez-Murillo et al., 2012). In addition to starch compositions, other parameters including physicochemical compositions are also important to determine the nutritional value of a given cereal. Yet for many product it has been shown that the ash content is often correlated with carbon and minerals contents (Monti et al., 2008).

Among the minerals only iron was significantly correlated with ash (p < 0.001). Indeed all monovalent ions (K⁺ and Na⁺) and divalent ions (Fe²⁺, Ca²⁺ and Zn²⁺) analyzed in millet are negatively correlated with total sugar content (and significantly for Na⁺ and K⁺ that is to say, p < 0.05). Conversely, among the divalent ions, only Mg²⁺ was positively correlated with the total sugar content. Likewise Fe²⁺ was significantly correlated with reducing sugars (p < 0.01) and significant correlated with Zn²⁺ to protein (p < 0.05). The salts are known for their influence on the rheological behavior of simple sugars. In a study performed in Sudan on millet, it has been shown that fermentation could improve the millet's protein digestibility and therefore its nutritional quality (Hag et al., 2002). Note that in addition to iron, ash content was significantly correlated with total sugar content.

Correlation between cultivars and nutritional parameters according and Principal Components Analysis.

The analysis of the dispersion of the different variables followed by the screening of all samples following the axes 1 and 2 (Figure 1 A and B) shows that millet samples can be divided into four distinct groups. We have chosen descriptors of iron-rich millets (group G1), carbohydrate-rich millets (group G2), and two less nutritious millets (groups G3 and G4).

The iron-rich group (G1) was composed of four cultivars such as CL, CE, CH and CI. In addition to iron, this group is characterized by cultivars with high levels in ash, zinc and reducing sugars. The cultivars of group G1 could be recommended for preventing anemia because of their high iron content. Also, the body requires iron for the synthesis of oxygen transport proteins, in particular hemoglobin and myoglobin, and for the formation of enzymes involved in electron transfer (McDowell, 2003). For the zinc, it plays the important roles (structural and functional) in many macromolecules and is required for enzymatic reactions. Zinc ions participate in all aspects of intermediary metabolism, transmission, and regulation of the expression of genetic information, storage, synthesis, and action of peptide hormones and structural maintenance of chromatin and biomembranes (Meunier et al., 2005).

The "carbohydrates" group (G2 group) is characterized by the pearl millet cultivars (CS, CO, CG, CJ, CP, CN, CQ, CF, CV, CS and CM) containing a high content of starch, amylose, total sugars, protein and dry matter. The cultivars of this group could be used in diet needing the high content in protein and sugar

sources specially the child's and adolescents. They are not recommended for diabetics and older people, but they can be important in starch industry. The other cultivars are less nutritious and are classified into G3 and G4 groups.




Figure 1 Factors dispersion plot according to the Principal Component Analysis (PCA) method (A); Projection of the pearl millet samples on the first and second component axes (B),

CONCLUSION

We show in this study that Benin has a diversity of cultivated pearl millets. The assessment of the nutritional value of 22 varieties of millets revealed that concentrations of major and minor nutrients like carbohydrate, proteins, sugars as well as minerals, varied widely. In general, we found no "all-in-one" variety capable of providing all potential health benefits attributed to pearl millets. In contrast, millet varieties under study were found to possess relatively low levels of proteins and sugars, but high levels of starch, amylose, Mg²⁺, Zn²⁺, Fe²⁺. Variations in the levels of those nutrients were dependent on both millet variety and site of production. Nutrient-rich millets could be used to improve nutritive value of other local varieties produced in Benin. Finally, this study supports evidence that millet could be recommended for preventing malnutrition or dietary deficiency in populations using millet as the basic staple food, particularly children and rural farming communities. Considering the high levels of starch observed here, further studies investigating all three types of starch, and thus millet digestibility, are needed.

Acknowledgments: Authors would like to thank the IFS for funding through an application No. AI3C/30956R gives to the first author.

REFERENCES

- Adebiyi, J.A., Obadina, A.O., Mulaba-Bafubandi, A.F., Adebo, O.A., Kayitesi, E., 2016. Effect of fermentation and malting on the microstructure and selected physicochemical properties of pearl millet (*Pennisetum glaucum*) flour and biscuit. *J. Cereal Sci.* 70, 132–139. <http://dx.doi:10.1016/j.jcs.2016.05.026>
- Akeredolu, I.A., Addo, A.A., Akeredolu, O.A., 2005. Clinical evaluation of pearl millet conophor weaning mix as supplementary food for Nigerian children. *Brazilian Arch. Biol. Technol.* 48, 531–536. <http://dx.doi:10.1590/S1516-89132005000500004>
- Almeida-Dominguez, H.D., Serna-Saldivar, S.O., Gomez, M.H., Rooney, L.W., 1993. Production and nutritional value of weaning foods from mixtures of pearl millet and cowpeas. *Cereal Chem.* 7, 14–18.
- Almeida-Dominguez, H.D., Serna-Saldivar, S.O., Rooney, L.W., 1991. Properties of new and commercial sorghum hybrid for use in alkaline-cooked foods. *Cereal Chem.* 68, 25–30.
- AOAC, 2000. Official Methods of Analysis of Association of Official Analytical Chemist, in: Association of Official Analytical Chemistry, Washington DC. pp. 4–15.
- Brigitte Michale-Dorea, D. Sauvart, 1989. Influence de la nature du concentré, céréales ou de betterave, sur la pulpe digestion chez les ruminants, *INRA Production Animale* ; 2 (4), 235-244.
- Chandrashekar, A., Kirleis, A.W., 1988. Influence of protein on starch gelatinization in sorghum. *Cereal Chem.* 65, 457–462.
- Chávez-Murillo, C.E., Méndez-Montevalvo, G., Wang, Y.J., Bello-Pérez, L.A., 2012. Starch of diverse Mexican rice cultivars: Physicochemical, structural, and nutritional features. *Starch/Staerke* 64, 745–756. <http://dx.doi:10.1002/star.201200016>
- Chung, H.J., Liu, Q., Lee, L., Wei, D., 2011. Relationship between the structure, physicochemical properties and in vitro digestibility of rice starches with different amylose contents. *Food Hydrocoll.* 25, 968–975. <http://dx.doi:10.1016/j.foodhyd.2010.09.011>
- Devi, P.B., Vijayabharathi, R., Sathyabama, S., Malleshi, N.G., Priyadarisini, V.B., 2014. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: A review. *J. Food Sci. Technol.* 51, 1021–1040. <http://dx.doi:10.1007/s13197-011-0584-9>
- Dykes, L., Rooney, L., 2007. Phenolic Compounds in Cereal Grains and Their Health Benefits. *Cereal Foods World* 52, 105–111. <http://dx.doi:10.1094/CFW-52-3-0105>
- FAO, 2015. Protection sociale et agriculture: Briser le cercle vicieux de la pauvreté rurale.
- Hag, M.E. El, Tinay, A.H. El, Yosif, N.E., 2002. Effect of fermentation on nutritive value and in vitro protein digestibility of pearl millet. *Food Chem.* 77, 193–196.
- Hoover, R., Swamidas, G., Kok, L.S., Vasanthan, T., 1996. Composition and Physicochemical Properties of Starch From pearl millet grains. *Food Chem.* 56, 355–367. <http://dx.doi:10.4314/bajopas.v5i1.12>
- Idem, N.U.A., Showemimu, F.S., 2004. Cereal crop of Nigeria pp. 16-34.
- Jain, R.K., Bal, S., 1997. Physical properties of pearl millet. *J. Agric. Eng. Res.* 66, 85–91.
- Kothari, S.L., Kumar, S., Vishnoi, R.K., Kothari, A., Watanabe, K.N., 2005. Applications of biotechnology for improvement of millet crops: Review of progress and future prospects. *Plant Biotechnol.* 22, 81–88. <http://dx.doi:10.5511/plantbiotechnology.22.81>
- Kouassi J.B., Massara C.C., Sess D.E., Tiahou G.G., Djohan F.Y., 2013. Détermination des teneurs en Magnésium, Potassium, Manganèse et Sodium de deux variétés de gombo. *Journal of Applied Biosciences* 67:5219 – 5227
- Laurent F., 1988. Utilisation du blé et des céréales dans la ration de vaches laitières. *Annales de Zootech.* 37 (2), 117-132
- Lopez-Rubio, B.A., A.A.K., S., Gidley, M.J., 2009. Resistant starch in vitro and in vivo: Factors determining yield, structure, and physiological relevance., in: S. Kasapis, I. T. Norton, & J. B. Ubbink (Eds.), *Modern Biopolymer Science*. San Diego, CA: Academic Press., p. 449–510.
- Macauley, H., Ramadjita, T., 2015. Les cultures céréalières : riz, maïs, millet, sorgho et blé. 31p.
- McDowell, L.R. (2003) *Minerals in Animal and Human Nutrition*. Elsevier Science, Amsterdam.
- McKeown, N.M., Meigs, J.B., Liu, S., Wilson, P.W., Jacques, P.F., 2002. Whole-grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study. *Am. J. Clin. Nutr.* 76, 390–398.
- Meunier, N., O'Connor, J.M., Maiani, G., Cashman, K.D., Secker, D.L. and Ferry, M. (2005) Importance of Zinc in the Elderly: The ZENITH Study. *European Journal of Clinical Nutrition*, 59, S1-S4.
- Modu, S., Laminu, H.H., F., S.A., 2010. Evaluation of the nutritional value of a composite meal prepared from Pearl millet (*Pennisetum Typhoideum*) and Cowpea (*Vigna Unguiculata*). *Bayero Journal Pure Appl. Sci.* 3, 164–168.
- Monti, A., Di Virgilio, N., Venturi, G., 2008. Mineral composition and ash content of six major energy crops. *Biomass and Bioenergy* 32, 216–223. <http://dx.doi:10.1016/j.biombioe.2007.09.012>
- Mouquet-Rivier, C., Icard-Vernière, C., Guyot, J.P., H, T., Rochette, I., Trèche, S., 2008. Consumption pattern, biochemical composition and nutritional value of fermented pearl millet gruels in Burkina Faso. *Int. J. Food Sci. Nutr.* 59, 716–729. <http://dx.doi:10.1080/09637480802206389>
- N'guessan, Y.D., Bedikou, M.E., Zoue, L.T., Goualie, B.G., Niamke, S.L., 2014. Physicochemical, nutritive and safety evaluation of local cereal flours sold in areas of the District of Abidjan-Côte d'Ivoire. *J. Appl. Biosci.* 83, 7579–7594. <http://dx.doi:10.4314/jab.v83i1.8>
- Obizoba, I.C., Atii, J.V., 1994. Evaluation of the effect of processing techniques on the nutrient and antinutrient contents of pearl millet (*Pennisetum glaucum*) seeds. *Plant Foods Hum. Nutr.* 45, 23–34.
- Oboh, G., Oladunmoye, M.K., 2007. Changes in the nutrient and anti-nutrient content of micro-fungi fermented cassava flour produced from low- and medium-cyanide variety of cassava tubers. *African J. Biotechnol.* 6, 2150–2157. <http://dx.doi:10.1177/026010600701800405>
- Ragaei, S., Sayed, A. El, Noaman, M., 2006. Antioxidant activity and nutrient composition of selected cereals for food use. *Food Chem.* 98, 32–38. <http://dx.doi:10.1016/j.foodchem.2005.04.039>
- Rooney L.W., Pflugfelder R.L., 1986. Factors affecting starch digestibility with special emphasis on sorghum and corn. *Journal of Animal Sciences*, 63, 1607–1623.
- Shweta Malik, 2015. Pearl millet-nutritional value and medicinal uses. *IJARIE*, vol-1 issue-3, p414-418.
- Songré-Ouattara, L.T., Mouquet-Rivier, C., Icard-Vernière, C., Humblot, C., Diawara, B., Guyot, J.P., 2008. Enzyme activities of lactic acid bacteria from a pearl millet fermented gruel (ben-saalga) of functional interest in nutrition. *Int. J. Food Microbiol.* 128, 395–400. <http://dx.doi:10.1016/j.ijfoodmicro.2008.09.004>
- Songré-Ouattara, L.T., Mouquet-Rivier, C., Icard-Vernière, C., Rochette, I., Diawara, B., Guyot, J.P., 2009. Potential of amylolytic lactic acid bacteria to replace the use of malt for partial starch hydrolysis to produce African fermented pearl millet gruel fortified with groundnut. *Int. J. Food Microbiol.* 130, 258–264. <http://dx.doi:10.1016/j.ijfoodmicro.2009.02.002>
- Subramanian, V., Jambunathan, R., 1982. Properties of sorghum grain and their relationship to roti quality, in: *International Crops Research Institute for the Semi Arid Tropics*. pp. 280–288.
- Taylor, J.R.N., Schober, T.J., Bean, S.R., 2006. Novel food and non-food uses for sorghum and millets. *J. Cereal Sci.* 44, 252–271. <http://dx.doi:10.1016/j.jcs.2006.06.009>
- Tou, E.H., Guyot, J.P., Mouquet-Rivier, C., Rochette, I., Counil, E., Traore, A.S., Treche, S., 2006. Study through surveys and fermentation kinetics of the traditional processing of pearl millet (*Pennisetum glaucum*) into ben-saalga, a fermented gruel from Burkina Faso. *Int. J. Food Microbiol.* 106, 52–60. <http://dx.doi:10.1016/j.ijfoodmicro.2005.05.010>
- Vanisha S. Nambiar, JJ Dhaduk, Neha Sareen, Tosha Shah and Rujuta Desai, 2011. Potential Functional Implications of Pearl millet (*Pennisetum glaucum*) in Health and Disease. *Journal of Applied Pharmaceutical Science* 01 (10): 62-67.
- Yadav, D.N., Anand, T., Navnidhi, Singh, A.K., 2014. Co-extrusion of pearl millet-why protein concentrate for expanded snacks. *Int. J. Food Sci. Technol.* 49, 840–846. <http://dx.doi:10.1111/ijfs.12373>
- Zhu, L.-J., Liu, Q.-Q., Wilson, J.D., Gu, M.-H., Shi, Y.-C., 2011. Digestibility and physicochemical properties of rice (*Oryza sativa* L.) flours and starches differing in amylose content. *Carbohydr. Polym.* 86, 1751–1759. <http://dx.doi:10.1016/j.carbpol.2011.07.017>